

Phenolic compounds from the lichen *Parmotrema cristiferum* (Taylor) Hale

Nguyen Thi Ngoc Van^{1,*}, Nguyen Mong Anh Nguyet¹, Tran Thai Bao¹, Nguyen Thi Le Thuy², Bui Thanh Tung², Pham Nguyen Kim Tuyen³, Nguyen Thi Hoai Thu⁴, Nguyen Thi Anh Tuyet⁵, Huynh Bui Linh Chi⁶, Nguyen Kim Phi Phung¹, Ngo Thi Thuy Duong¹



Use your smartphone to scan this QR code and download this article

¹Department of Organic Chemistry, University of Science, National University - Ho Chi Minh City, Vietnam

²Department of Biotechnology, Ho Chi Minh City Open University, Vietnam

³Faculty of Environmental Science, Sai Gon University, Ho Chi Minh City

⁴Faculty of Basic Sciences, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Vietnam

⁵Department of Chemistry, Ho Chi Minh City University of Education, Ho Chi Minh City, Vietnam

⁶Department of Science, Dong Nai University, Dong Nai Province

Correspondence

Nguyen Thi Ngoc Van, Department of Organic Chemistry, University of Science, National University - Ho Chi Minh City, Vietnam

Email: ntnvan101096@gmail.com

History

- Received: 2021-08-08
- Accepted: 2021-09-10
- Published: 2021-09-21

DOI: 10.32508/stdj.v24i4.3781



Copyright

© VNU-HCM Press. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



ABSTRACT

Introduction: Lichens are fungal and algal or cyanobacterial symbioses resulting in the production of specific metabolites with a great variety of effects such as antimycobacterial, antiviral, anti-inflammatory, antipyretic and antiproliferative. *Parmotrema cristiferum* (Taylor) Hale, a species of foliose lichen, was collected at Champasack Province, Laos in April 2015. This paper describes the isolation and structure elucidation of nine compounds isolated from this lichen. **Methods:** Phytochemical investigations of the chloroform extract of the lichen *P. cristiferum* led to the isolation of nine pure compounds. Their chemical structures were elucidated by extensive HR-ESI-MS and 1D and 2D-NMR spectroscopic analysis, and comparison with previously published data. **Results:** From the studied lichen, nine compounds, orcinol (**1**), orsellinic acid (**2**), atranol (**3**), methyl β -orsellinate (**4**), atranorin (**5**), diffractaic acid (**6**), lecanoric acid (**7**), baileisidone (**8**), and tinctorinone (**9**) were isolated, structurally elucidated and evaluated for their α -glucosidase inhibitory activity. **Conclusions:** Among isolated compounds, except **2**, **4**, and **7**, the remaining six compounds were known for the first time from the lichen *P. cristiferum*. All these compounds were inactive in the α -glucosidase enzyme inhibitory assay.

Key words: *Parmotrema cristiferum* (Taylor) Hale, depside, depsidone, phenolic, α -glucosidase

INTRODUCTION

Diabetes mellitus is a metabolic disorder characterized by chronic hyperglycemia. There are many and diverse therapeutic strategies in the management of type 2 diabetes. Predominantly herbal drugs have been widely used globally for diabetic treatment over thousands of years due to their traditional acceptability and lesser side effects. Therefore, screening of α -amylase and α -glucosidase inhibitors in medicinal plants has received much attention.¹

Parmotrema cristiferum (Taylor) Hale is a foliose lichen; belonging to the genus *Parmotrema*, one of the largest genera of the Parmeliaceae family.² Previous studies of this lichen reported that its methanolic extract had various pharmacological activities such as antimicrobial, antioxidant, and insecticidal activities.³ However, metabolites possessing antidiabetic activity unexplored well as no chemical data of this lichen are reported. Therefore, to search for new classes of α -glucosidase inhibitors, we nine compounds' isolation and structural elucidation from the chloroform extract of the lichen *Parmotrema cristiferum* (Taylor) Hale and the α -glucosidase inhibitory activity of these compounds.

MATERIALS AND METHODS

General experimental procedures

The HR-ESI-MS was recorded on an HR-ESI-MS MicroTOF-Q mass spectrometer. The ¹H-NMR (500 MHz) and ¹³C-NMR (125 MHz) spectra were recorded on a Bruker Avance 500 spectrometer. Chemical shifts are expressed in ppm using a residual solvent signal as internal reference (CDCl₃ δ_H 7.26, δ_C 77.2). α -Glucosidase (EC 3.2.1.20) from *Saccharomyces cerevisiae* (750 UN) and *p*-nitrophenyl- α -D-glucopyranoside were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Acarbose and dimethyl sulfoxide were obtained from Merck. Other chemicals were of the highest grade available.

Thin-layer chromatography (TLC) was carried out on precoated silica gel 60 F254 or silica gel 60 RP-18 F254S (Merck) and the isolated compounds were visualized by spraying with 5% vanillin solution followed by heating. Gravity column chromatography was performed on silica gel 60 (0.040–0.063 mm, Himedia).

Cite this article : Van N T N, Nguyet N M A, Bao T T, Thuy N T L, Tung B T, Tuyen P N K, Thu N T H, Tuyet N T A, Chi H B L, Phung N K P, Duong N T T. **Phenolic compounds from the lichen *Parmotrema cristiferum* (Taylor) Hale.** *Sci. Tech. Dev. J.*; 24(4):2134-2140.

Plant material

The lichen thalli of *Parmotrema cristiferum* (Taylor) Hale was collected at 109 m altitude at Paksong town (15°11'00"N, 106°14'00"E), Paksong district, Champasack province, Laos in April 2015. The botanical species of *Parmotrema cristiferum* (Taylor) Hale was authenticated by Dr. Vo Thi Phi Giao, Faculty of Biology, University of Science, National University — Ho Chi Minh City.

Extraction and isolation

The thallus material (1.2 kg) was washed underflow of tap water and then was air-dried at ambient temp to obviate thermally induced decomposition before being ground into a fine powder. The ground powder sample (0.8 kg) was macerated by acetone at room temperature to afford a crude acetone extract (79.0 g). This crude was subjected to silica gel solid-phase extraction and eluted consecutively with *n*-hexane, chloroform, ethyl acetate, acetone, and methanol at room temperature and then each type of eluants was evaporated at reduced pressure to afford *n*-hexane (6.39 g), chloroform (30.48 g), ethyl acetate (32.57 g), acetone (7.82 g), and methanol (1.64 g) extracts, respectively. The chloroform extract (30.48 g) was applied to silica gel column chromatography and eluted with solvent systems of *n*-hexane:chloroform (stepwise, 9:1 to 0:10) to give five fractions, coded C1 (1.18 g), C2 (11.59 g), C3 (15.49 g), C4 (1.10 g), C5 (1.12 g). Fraction C1 (1.18 g) was repeatedly chromatographed followed by preparative TLC, eluted with *n*-hexane:chloroform (5:5 to 0:10) to afford four compounds coded 1 (18.0 mg), 2 (7.2 mg), 3 (12.3 mg), and 4 (64.0 mg). Fraction C2 (11.59 g) was silica gel rechromatographed and eluted with chloroform:ethyl acetate (6:4 to 0:10) to give three compounds: 5 (95.3 mg), 6 (13.8 mg), and 7 (7.3 mg). Fraction C3 (15.49 g) was selected for further fractionation by silica gel CC using an isocratic mobile phase consisting of chloroform:methanol (9:1 to 5:5) to obtain six subfractions (C3.1 - C3.6). Further fractionation of C3.2 (586.0 mg) was applied to a silica gel CC using chloroform:methanol (98:2) to afford 8 (10.0 mg), and 9 (5.3 mg).

α -Glucosidase inhibition assay

The inhibitory activity of α -glucosidase was determined as our previous assay.⁴ Reaction mixture consisting of 60 mL of phosphate-buffered solution (100 mM, pH 6.8), 20 mL of test samples at different concentrations, and 20 mL of α -glucosidase (0.3 IU.mL⁻¹) was injected into a 96 well plate and incubated at 37 °C for 10 minutes. Then, 100 mL

of *p*-nitrophenyl- α -D-glucopyranoside (200 mM) solution was added to this mixture in a phosphate buffer. Next, incubate the reaction mixture at 37 °C for 30 minutes. Finally, add 50 mL of NaOH solution (50 mM) to this mixture. The absorbance of *p*-nitrophenol (*p*NP) was measured at 405 nm on the ELX800 (BIOTEX). Acarbose was used as the positive control.

RESULTS

By using efficient separation techniques, the chemical investigation on the chloroform extract of the lichen *P. cristiferum* led to the isolation of nine compounds. Their chemical structures were elucidated by 1D and 2D NMR as well as the HR-ESI-MS. They were four phenolic compounds, orcinol (1), orsellinic acid (2), atranol (3), methyl β -orsellinate (4), three depsides, atranorin (5), diffractaic acid (6), lecanoric acid (7), and two depsidones, baileisidone (8), and tinctorinone (9). Their ¹H and ¹³C-NMR data of compounds 1-4 were presented in Tables 1 and 3; of compounds 5-9 were presented in Tables 2 and 3; and the following characteristics.

- Orcinol (1): Colorless needles, mp. 106-108 °C. HR-ESI-MS (positive mode) *m/z* 125.0601 [M+H]⁺ (calcd. for C₇H₈O₂ 125.0602).
- Orsellinic acid (2): Colorless needles, mp. 184-185 °C. HR-ESI-MS (positive mode) *m/z* 169.0491 [M+H]⁺ (calcd. for C₈H₈O₄+H, 169.0501).
- Atranol (3): Faint yellow needles, mp. 124 °C. HR-ESI-MS (positive mode) *m/z* 153.0536 [M+H]⁺ (calcd. for C₈H₈O₃+H, 153.0551).
- Methyl β -orsellinate (4): Colorless needles, mp. 143-144 °C. HR-ESI-MS (positive mode) *m/z* 197.0814 [M+H]⁺ (calcd. for C₁₀H₁₂O₄+H, 197.0811).
- Atranorin (5): Prisms, mp. 196 °C. HR-ESI-MS (negative mode) *m/z* 373.0881 [M-H]⁻ (calcd. for C₁₉H₁₈O₈-H, 373.0923).
- Diffractaic acid (6): White amorphous powder, mp. 189-190 °C. HR-ESI-MS (positive mode) *m/z* 375.1437 [M+H]⁺ (calcd. for C₂₀H₂₂O₇+H, 375.1443).
- Lecanoric acid (7): Colorless needles, mp. 184 °C. HR-ESI-MS (negative mode) *m/z* 317.0672 [M-H]⁻ (calcd. for C₁₆H₁₄O₇-H, 317.0661).
- Baileisidone (8): White amorphous powder, $[\alpha]_D^{29} = 0$ {[α]_D²⁵ 47.4 (c 0.7, CHCl₃)}.⁵ HR-ESI-MS (positive mode) *m/z* 415.1025 [M+H]⁺ (calcd. for C₂₁H₁₈O₉+H, 415.1029).

- Tinctorinone (**9**): White amorphous powder. $[\alpha]_D^{29} = 0$. HR-ESI-MS (positive mode) m/z 415.1025 $[M+H]^+$ (calcd. for $C_{21}H_{18}O_9+H$, 415.1029).

DISCUSSION

The structures of nine isolated compounds were elucidated based on modern physicochemical methods such as HR-MS and 1D and 2D NMR compared their data with those in the literature. The compounds possessing a similar framework were structurally discussed.

Compound **1** was isolated as colorless needles. Its molecular formula was determined as $C_7H_8O_2$ through its molecular ion peak at m/z 125.0601 $[M+H]^+$ in the HR-ESI-MS spectrum. The 1H -NMR data of **1** gave one methyl group, three aromatic methine protons, and two hydroxyl protons (Table 1). The ^{13}C -NMR spectrum showed five signals, including one methyl group, three aromatic methines, and three quaternary aromatic carbons (Table 3). The good compatibility of its NMR and MS data with those of orcinol in the literature^{6,7} proposed that compound **1** was orcinol.

Compound **2** was obtained as colorless needles. The similarity in the NMR data (Tables 1 and 3) just with the replacement of one aromatic proton in the **1** by a carboxyl group in **2** suggested that it was orsellinic acid.^{6,7} This was supported by the pseudomolecular ion peak at m/z 169.0491 $[M+H]^+$ in the HR-ESI-MS spectrum which fitted well with the molecular formula of $C_8H_8O_4$. Therefore, **2** was orsellinic acid.

Compound **3** was quickly identified as atranol because it possessed similar NMR data (Tables 1 and 3) with those in the literature.^{6,8} The position of the aldehyde group was clarified by HMBC cross-peaks of its proton (δ_H 10.23) to C-3 (δ_C 109.3) and C-2, 4 (δ_C 163.2) (Figure 2). Its molecular formula $C_8H_8O_3$ was determined through the pseudomolecular ion peak at m/z 153.0536 $[M+H]^+$.

The comparison NMR data of **1** and **4** showed that the latter possessed one more methoxycarbonyl (δ_H 3.71, δ_C 173.5, 52.1, COO-CH₃) and one methyl group (δ_H 1.82, δ_C 8.1, C-9). The positions of these two groups were evidenced by HMBC cross-peaks of the methoxy protons at δ_H 3.71 (7-OCH₃) with the carboxyl carbon at δ_C 173.5 (C-7), and of the methyl protons (δ_H 1.82) with adjacent aromatic carbons at δ_C 164.1 (C-2), 109.5 (C-3) and 161.0 (C-4). Based on the compatibility of the NMR and HR-ESI-MS data of **4** with the literature,⁹ **4** was methyl β -orsellinate.

Compound **5** was preliminarily proposed as a depside via the comparison of its NMR data (Tables 2 and 3) with corresponding depsides in the literature. A close search showed that **5** was composed of the combination of **3** and **4** moieties. This observation was well supported by its HMBC pattern (Figure 1) and HR-ESI-MS spectrum (negative mode) with a pseudomolecular ion peak at m/z 373.0881 $[M-H]^-$ (calcd. for $C_{19}H_{18}O_8-H$, 373.0923). Thus, the structure of **5** was assigned as atranorin.⁸

Compound **6** was obtained as a white amorphous powder. A careful search of the NMR data (Tables 2 and 3) showed that **6** was also a depside and composed of two moieties of β -orsellinic acid (the carboxylic acid of **4**). This was proved by the HR-ESI-MS spectrum (positive mode) with a pseudomolecular ion peak at m/z 375.1437 $[M+H]^+$ (calcd. for $C_{20}H_{22}O_7+H$, 375.1443), by HMBC experiment (Figure 1), and by the similar NMR data with those of diffractaic acid.^{10,11} Finally, the chemical structure of **6** was determined to be diffractaic acid.

Compound **7** was isolated as colorless needles. In the same above argument, the similarity in the spectral data (Tables 2 and 3) suggested that **7** was composed of two moieties of orsellinic acid (**2**). The HR-MS spectrum (m/z 317.0672 $[M-H]^-$, calcd. for $C_{16}H_{14}O_7-H$, 317.0661) and good accordance in the NMR data¹²(Tables 1 and 2) proposed **7** as lecanoric acid.

Compound **9** was isolated as a white amorphous powder. The comparison NMR data of **9** and **8** (Tables 2 and 3) showed that they were depsidones with a lactone ring, with some differences. HMBC experiment of **9** confirmed that its lactone moiety possessed one acetal carbon (δ_H 7.10, δ_C 102.4) and one ethoxy side-chain [δ_H 3.97, δ_C 66.2 (O-CH₂-CH₃), and δ_H 1.34 (t, 7.0 Hz), δ_C 15.6 (O-CH₂-CH₃)]. In addition, its A ring contained a further hydroxymethylene [δ_H 4.96 (dd, 11.5, 8.5 Hz), 4.79 (dd, 11.5, 8.5 Hz), δ_C 53.0, C-9]. The comparison of these NMR data with those of tinctorinone, a depsidone in the lichen *Parmotrema tinctorum*¹³ showed the compatibility. The specific optical rotation data of **9** was zero, thus it was a racemic mixture of tinctorinone.

The inhibitory assay against α -glucosidase for isolated compounds was evaluated. The results (Table 4) indicated that these compounds (1-9) were inactive.

CONCLUSION

From the chloroform extract of the lichen *Parmotrema cristiferum*, collected at Paksong district,




Table 1: ¹H-NMR (Acetone-d₆) data of compound 1-4

No.	1	2	3	4
1	6.15 s		6.23 s	
3	6.15 s	6.29 (d, 2.0)		
5	6.15 s	6.22 (d, 2.5)	6.23 s	6.14 s
7	2.15 s		2.20 s	
8		2.51 s	10.23 s	2.21 s
9				1.82 s
COOCH ₃				3.71 s
2-OH	8.02 s		10.85 s	11.73 s
4-OH	8.02 s		10.85 s	9.02 s




Table 2: ¹H-NMR data of compound 5-9

No.	5 ^(b)	6 ^(a)	7 ^(a)	8 ^(c)	9 ^(a)
3			6.14 (d, 2.5)		
5	6.42 s	6.76 s	6.23 (d, 1.5)	6.60 s	6.95 s
8	10.21 s	2.45 s	2.44 s	2.44 s	2.49 s
9a	2.39 s	2.14 s			4.96 (dd, 11.5, 8.5)
9b					4.79 (dd, 11.5, 4.5)
2-OCH ₃		3.85 s			
4-OCH ₃		3.91 s		3.92 s	3.94 s
2-OH			11.10 s		
9-OH					4.16 (dd, 8.5, 4.5)
3'			6.44 (d, 2.5)		
5'	6.65 s	6.57 s	6.36 (d, 2.5)		
8'	2.04 s	2.13 s	2.49 s	2.26 s	2.25 s
9'	2.34 s	2.64 s		6.15 (dd, 8.5, 3.0)	7.10 s
10'a				3.05 (dd, 17.5, 8.5)	4.04 m
10'b				3.61 (dd, 17.5, 3.0)	3.97 m
11'					1.34 (t, 7.0)
12'				2.29 s	
7'-OCH ₃	3.88 s				
2'-OH	10.51 s				

(a) Acetone-*d*₆; (b) DMSO-*d*₆; (c) CDCl₃

Champasack province, Laos, using various chromatographic methods, nine phenolic compounds were isolated. They were orcinol (1), orsellinic acid (2), atranol (3), methyl β-orsellinate (4), atranorin (5), diffractaic acid (6), lecanoric acid (7), a racemic mixture of baileisidone (8), and a racemic mixture of tinctorinone (9). Except for compounds 2, 4, and 7,¹⁴ the remaining six compounds were reported for the first time in this species. All nine compounds were evaluated against α-glucosidase enzyme but none of them were active. Further research on the remaining extracts is continued.

ABBREVIATIONS

HR-ESI-MS: High resolution- Electrospray ionization-Mass spectrometry

¹H-NMR: Proton Nuclear Magnetic Resonance

¹³C-NMR: Carbon-13 Nuclear Magnetic Resonance

HMBC: Heteronuclear Multiple Bond Correlation

s: singlet

d: doublet

dd: doublet of doublets

t: triplet

m: multiplet

COMPETING INTEREST

The authors declare no competing financial interest.

AUTHORS' CONTRIBUTION

Huynh B.L.C, Ngo T.T.D contributed in conducting experiments, acquisition of data, interpretation of data. Nguyen T.N.V, Nguyen M.A.N, Tran T.B interpreted NMR and MS data as well as searched the bibliography. Nguyen T.L.T, Bui T. T did the biological assay. Nguyen K.P.P. gave the final correction for the manuscript

Corresponding author: Nguyen Thi Ngoc Van, University of Science, National University-Ho Chi Minh

Table 3: ¹³C-NMR data of compound 1-9

No.	1 ^(a)	2 ^(a)	3 ^(a)	4 ^(a)	5 ^(b)	6 ^(a)	7 ^(a)	8 ^(c)	9 ^(a)
1	108.4	104.8	108.4	104.9	110.7	121.3	104.9	114.3	114.2
2	159.4	163.4	163.2	164.1	163.5	157.8	166.6	147.6	160.1
3	100.7	101.5	109.3	109.5	107.9	117.8	101.8	134.4	119.5
4	159.4	167.1	163.2	161.0	161.6	160.8	159.1	150.9	162.9
5	108.4	112.1	108.4	111.5	109.0	109.1	112.8	110.9	112.3
6	140.6	144.9	151.6	140.5	149.0	135.9	144.6	134.4	145.9
7	21.5	174.2	22.4	173.5	164.6	166.6	167.9	161.5	162.0
8		24.2	194.3	24.2	193.9	20.0	24.3	20.7	21.4
9				8.1	21.2	9.0			53.0
2-OCH ₃						62.3			
4-OCH ₃						56.2		56.6	56.7
COOCH ₃				52.1					
1'					115.3	113.1	116.2	107.2	109.1
2'					157.4	163.9	165.3	152.5	152.8
3'					116.3	116.8	108.4	119.1	121.3
4'					151.4	153.3	153.0	149.9	150.4
5'					115.7	116.1	115.5	137.9	139.7
6'					136.6	141.5	144.6	136.2	133.9
7'					169.7	175.4	170.6	170.9	169.1
8'					9.3	9.4	23.9	9.1	9.3
9'					21.1	23.9		76.9	102.4
10'								46.7	66.2
11'								204.3	15.6
12'								30.7	
7'-OCH ₃					52.3				

(a) Acetone-*d*₆; (b) DMSO-*d*₆; (c) CDCl₃

City, 227 Nguyen Van Cu Street, Ward 4, District 5, Ho Chi Minh City. Email: ntnvan101096@gmail.com

ACKNOWLEDGMENT

We would like to thank Dr. Vo Thi Phi Giao for the lichen authentication.

REFERENCES

- Funke I, Melzig MF. Traditionally used plants in diabetes therapy: phytotherapeutics as inhibitors of alpha-amylase activity. *Revista Brasileira de Farmacognosia*, 16 (1), 1-5, 2006; Available from: <https://doi.org/10.1590/S0102-695X2006000100002>.

- Ahmadjian V, Hale ME. *The Lichens*. Academic Press, New York London, 1973;.
- Kekuda T, et al. Radical scavenging, antimicrobial and insecticidal efficacy of *Parmotrema cristiferum* and *Dirinaria applanata*. *Science, Technology and Arts Research Journal*, 4 (1), 95, 2015; Available from: <https://doi.org/10.4314/star.v4i1.16>.
- Nguyen HP, et al. A new glycoside and in vitro evaluation of α -Glucosidase inhibitory activity of constituents of the mangrove *Lumnitzera racemosa*. *Natural Product Communications*, 12 (11), 1751-1754, 2017; Available from: <https://doi.org/10.1177/1934578X1701201125>.
- Nguyen VK, et al. Chemical constituents of the lichen *Usnea baileyi* (Stirt.) Zahlbr. *Tetrahedron Lett.*, vol. 59 (14), 1348-1351, 2018; Available from: <https://doi.org/10.1016/j.tetlet.2018.04.011>.

Table 4: α -Glucosidase inhibitory activity of isolated compounds

No.	Compound and coded number	IC ₅₀ (μ M) \pm SD
1	Orcinol (1)	1523.5 \pm 4.0
2	Orselinic acid (2)	441.0 \pm 3.0
3	Atranol (3)	599.2 \pm 2.6
4	Methyl β -orselinate (4)	360.0 \pm 1.5
5	Atranorin (5)	262.8 \pm 1.0
6	Diffractaic acid (6)	419.6 \pm 0.5
7	Lecanoric acid (7)	579.7 \pm 1.6
8	Bailesidone (8)	234.5 \pm 0.7
9	Tinctorinone (9)	304.2 \pm 0.7
10	Acarbose (Positive control)	209.7 \pm 0.3

- tetlet.2018.02.007.
- Huneck S, et al. Identification of lichen substances. Springer. New York, 1996; Available from: <https://doi.org/10.1007/978-3-642-85243-5>.
 - Lopes TIB, et al. Radical-scavenging activity of orsellinates. Chemical and Pharmaceutical Bulletin, 56 (11), 1551-1554, 2008; PMID: 18981604. Available from: <https://doi.org/10.1248/cpb.56.1551>.
 - Nguyen TTT, et al. Study on chemical constituents of the lichen Parmotrema tinctorum (Nyl.) Hale. Vietnam Journal of Science and Technology, 56 (4), 434, 2018; Available from: <https://doi.org/10.15625/2525-2518/56/4/11638>.
 - Hylands PJ, Ingolfsdottir K. The isolation of methyl β -orsellinate from Stereocaulon alpinum and comments on the isolation of 4, 6-dihydroxy-2-methoxy-3-methylacetophenone from Stereocaulon species. Phytochemistry, 24 (1), 127-129, 1985; Available from: [https://doi.org/10.1016/S0031-9422\(00\)80821-3](https://doi.org/10.1016/S0031-9422(00)80821-3).
 - Brandão LFG, et al. Cytotoxic evaluation of phenolic compounds from lichens against melanoma cells. Chemical and Pharmaceutical Bulletin, 61 (2), 176-183, 2013; PMID: 23207680. Available from: <https://doi.org/10.1248/cpb.c12-00739>.
 - Nishitoba Y, et al. Lichen acids, plant growth inhibitors from Usnea longissima. Phytochemistry, 26 (12), 3181-3185, 1987; Available from: [https://doi.org/10.1016/S0031-9422\(00\)82466-8](https://doi.org/10.1016/S0031-9422(00)82466-8).
 - Choudhary MI, et al. New antiglycation and enzyme inhibitors from Parmotrema cooperi. Science China Chemistry, 54 (12), 1926-1931, 2011; Available from: <https://doi.org/10.1007/s11426-011-4436-2>.
 - Bui VM, et al. Two new phenolic compounds from the Vietnamese lichen Parmotrema tinctorum. Natural Product Research; Available from: <https://doi.org/10.1080/14786419.2020.1864367>.
 - Pham NKT, et al. Two new phenolic compounds from the lichen Parmotrema cristiferum growing in Vietnam. Natural Product Research, 1-7; Available from: <https://doi.org/10.1080/14786419.2021.1892672>.